

## Harmonic Analysis Homework Sheet 3

### Exercise 3.1

Establish Theorem 1.3.4 in the case where one of the  $q_i \leq 1$ . The following lemma might be helpful.

**Lemma 0.1.** *Let  $\Lambda_\lambda(x, y) = (1 - \lambda)x + \lambda y$  for  $0 \leq \lambda \leq 1$  and  $x, y > 0$ . Then,*

$$\Lambda_\alpha(\Lambda_\beta(x, y), \Lambda_\gamma(x, y)) = \Lambda_{\Lambda_\alpha(\beta, \gamma)}(x, y).$$

### Exercise 3.2

Let  $A, B_0, B_1 \geq 0$  such that  $A \leq \min(B_0, B_1)$ . Show that we have

$$A \leq B_\theta \min\left(\frac{B_0}{B_1}, \frac{B_1}{B_0}\right)^\varepsilon$$

for any sufficiently small  $\varepsilon > 0$ , where  $B_\theta = B_0^{1-\theta} B_1^\theta$ . Can you imagine a situation where this type of estimate is superior to the simpler estimate  $A \leq B_\theta$ ?

### Exercise 3.3

In Lemma 1.3.5, show that the sub-double exponential hypothesis  $|f(z)| \lesssim_f \exp(\mathcal{O}_f(e^{(\pi-\delta)|z|}))$  for some  $\delta > 0$  is completely sharp.

### Exercise 3.4

Prove Hadamard's three circle theorem. Let  $g(z)$  be holomorphic on the annulus  $A := \{z \in \mathbb{C} : r_1 \leq |z| \leq r_3\}$  for some  $0 < r_1 < r_3$  and denote

$$M(r) := \max_{\theta \in [0, 2\pi]} |g(re^{i\theta})| \quad \text{for } r \in (r_1, r_3).$$

Prove that

$$\log\left(\frac{r_3}{r_1}\right) \log M(r_2) \leq \log\left(\frac{r_3}{r_2}\right) \log M(r_1) + \log\left(\frac{r_2}{r_1}\right) \log M(r_3)$$

for any  $r_1 < r_2 < r_3$ , i.e.,  $\log M(r)$  is a convex function of  $\log r$ . (Hint: Convince yourself that there is a vertical strip which the exponential function maps onto the annulus  $A$ .)

Compute both sides of the inequality when  $g(z) = cz^\lambda$  for some constants  $c \in \mathbb{C}$  and  $\lambda \in \mathbb{Z}$ .