

Harmonic Analysis Homework Sheet 8

Exercise 8.1

Compute the Dirichlet kernel, i.e., for $t \in [0, 1]$ and $N \in \mathbb{N}$ show that

$$D_N(t) := \sum_{m=-N}^N e^{2\pi i m t} = \frac{\sin((2N+1)\pi t)}{\sin(\pi t)}.$$

Note that the higher-dimensional *square Dirichlet kernel*, i.e.,

$$D_N^\square(t) := \sum_{m \in \mathbb{Z}^d, |m_j| \leq N} e^{2\pi i m \cdot t} = \prod_{j=1}^d D_N(t_j), \quad t \in \mathbb{T}^d$$

simply factorizes (as opposed to the *spherical Dirichlet kernel* $\sum_{m \in \mathbb{Z}^d, |m| \leq N} e^{2\pi i m \cdot t}$).

Exercise 8.2

The disc conjecture for $1 < p < \infty$ is the statement that the Fourier multiplier $\mathbf{1}_{B_0(1)}(\xi)$ is bounded on $L^p(\mathbb{R}^2)$, i.e., $\|(\mathbf{1}_{B_0(1)} \hat{f})^\vee\|_p \lesssim_p \|f\|_p$. (The case $p = 2$ is trivial in view of Plancherel's theorem.) Show the following

Lemma 0.1 (Y. Meyer). *Let $(v_j)_{j \in \mathbb{N}} \in \mathbb{S}^1$ be a sequence of unit vectors in \mathbb{R}^2 and let H_j be the half-plane $\{x \in \mathbb{R}^2 : x \cdot v_j \geq 0\}$. Define the “half plane multipliers” $(T_j)_{j \in \mathbb{N}}$ on $L^p(\mathbb{R}^2)$ by setting $\widehat{T_j f}(\xi) = \mathbf{1}_{H_j}(\xi) \hat{f}(\xi)$. If the disc conjecture holds for some $1 < p < \infty$, then for any sequence $(f_j)_{j \in \mathbb{N}}$, we have the square function estimate*

$$\left\| \left(\sum_j |T_j f_j|^2 \right)^{1/2} \right\|_p \lesssim_p \left\| \left(\sum_j |f_j|^2 \right)^{1/2} \right\|_p, \quad f_j \in \mathcal{S}(\mathbb{R}^2).$$

Exercise 8.3

Truncated integral kernels usually give rise to somewhat “localized” operators. For $R > 0$, we say that a linear operator T is R -local if $\text{supp } Tf \subseteq R \text{supp } f$. (Here, $RA = \{x \in \mathbb{R}^d : x \in B_y(R) \text{ for all } y \in A\}$ for some $A \subseteq \mathbb{R}^d$.) In particular, this means that $\text{supp } T\mathbf{1}_{B_x(R)} \subseteq B_x(2R)$. Show the following

Lemma 0.2. *Suppose T is a R -local linear operator taking functions on \mathbb{R}^d to functions on \mathbb{R}^d . Then, for any $1 \leq p \leq q \leq \infty$, the bound*

$$\|Tf\|_q \lesssim \|f\|_p, \quad f \in L^p(\mathbb{R}^d) \tag{1}$$

is equivalent to the bound

$$\|Tf\|_{L^q(B_x(2R))} \lesssim \|f\|_p, \quad f \in L^p(B_x(R)) \tag{2}$$

holding uniformly in $x \in \mathbb{R}^d$. In other words, to show (1) it suffices to test it for functions on an R -ball.