

# Harmonic Analysis (Winter term 2023/2024)

October 2, 2023

**Lectures** (four hours per week): ??? ??-??-?? (in ????) and ??? ??-??-?? (in ????), starting on October ??, 2023

**Exercises** (two hours per week): ??? ??-??-?? (in ????)

**Credits:** 10

**Prerequisites:** Analysis 1-3.

Analysis in general tends to revolve around the study of general classes of *functions* and *operators*. *Real-variable harmonic analysis* focuses in particular on the relation between *qualitative* properties (such as measurability, boundedness, differentiability, analyticity, integrability, decay at infinity, convergence, etc.) and their *quantification* (i.e., what is the smallest upper bound on a function, how often is it differentiable, what is its  $L^p$  norm, what is the convergence rate of a sequence, etc.). It is then natural to ask how quantitative properties of such functions change when one applies various (often quite explicit) operators. It turns out that quantitative estimates, such as  $L^p$  estimates on such operators, provide an important route to establish qualitative results and in fact there are a number of principles (such as the uniform boundedness principle or Stein's maximal principle [21]) which assert that this is the *only* route, in the sense that a quantitative result must exist in order for the qualitative result to be true.

Many arguments in harmonic analysis will, at some point, involve a combinatorial statement about certain types of geometric objects such as cubes, balls, or tubes. One such useful statement is the *Vitali* covering lemma which asserts that given a collection of balls  $B_1, \dots, B_k$  in  $\mathbb{R}^d$ , then there exists a subcollection of balls  $B_{i_1}, \dots, B_{i_m}$  which are disjoint but contain a significant fraction of the volume covered by the original balls, in the sense that  $|\bigcup_{i=1}^k B_i| \leq a_d |\bigcup_{j=1}^m B_{i_j}|$  for some  $d$ -dependent constant  $a_d$ .

One feature of harmonic analysis methods is that they tend to be *local* rather than *global*. For instance, it is quite common to analyze a function  $f$  by applying cutoff functions in either the spatial or the frequency variables to *decompose*  $f$  into a number of somewhat localized pieces. One then estimates each of these pieces separately and “glues” the estimates back together at the end. One reason for this “divide and conquer” strategy is that generic functions tend to have infinitely many degrees of freedom ( $f$  may for instance be very smooth but slowly decaying at one place whereas at other places  $f$  may be highly singular

or oscillating very quickly) and it would be quite difficult to treat all of these features at once. A well chosen decomposition can isolate these features from each other, so that each component only has one salient feature that could cause difficulty. In reassembling the estimates from the individual components, one can use rather crude tools such as the triangle inequality, or more refined tools, such as ones relying on *(almost) orthogonality*. The main drawback of decomposition methods is however that one generally does not obtain the optimal constants.

Another basic theme of harmonic analysis is the attempt to quantify the elusive phenomenon of *oscillation*. Intuitively, if an expression oscillates wildly in phase, then its average value should be relatively small in amplitude. This leads to the *principle of stationary phase* and the *Heisenberg uncertainty principle* which relates the decay and smoothness of a function to the smoothness and the decay of its Fourier transform. The development of a robust theory for *oscillatory integrals* is also one main ingredient to understand the interplay between  $L^p$  estimates of certain Fourier multipliers (such as the *Bochner–Riesz means*) and geometric properties of certain (smooth) manifolds, such as the decay of the Fourier transform of the associated surface measure.

1. Lebesgue spaces, Lorentz spaces, interpolation (Riesz–Thorin, Marcinkiewicz (in particular restricted weak-type formulation. See, e.g., Tao [26, Lecture 2] or Stein [25]).
2. Covering lemmas (Vitali, Whitney, Calderón–Zygmund) and Calderón–Zygmund decomposition. See, e.g., Stein [22, Chapter 1], Grafakos [14, Proposition 2.1.20, Theorem 4.3.1] and Guzmán [9].
3. Maximal functions. See Stein [22, 24].
  - (a) Hardy–Littlewood maximal function and Lebesgue differentiation theorem. See Stein [22, Chapter 1] and [24, Chapter I, Section §3].
  - (b) Maximal functions and Lebesgue differentiation for more general sets (instead of balls). See Guzmán [9] and Cordoba–Fefferman [8].
  - (c) Hardy–Littlewood  $p$  maximal function. See Blunck–Kunstmann [3, 1].
  - (d) Relation to convergence almost everywhere, first glance at Bochner–Riesz summability (FAP1 and FAP2 in [17]. In this regard, see also the “ergodic Hopf–Dunford–Schwartz” theorem [23, p. 48] respectively Dunford–Schwartz [10] (Section XIII.6: Lemma 7 (p. 676), Theorem 8 (p. 678); Section XIII.8: Lemma 6 (p. 690) Theorem 7 (p. 693); Section XIII.9: Exercise 3 (p. 717))).
4. Singular integrals following Stein [22, Chapter II], [24, Chapter I, Section §5]. In particular, Hilbert transform and its application to partial sums operators [22, Chapter IV, Section §4] and second glance at Bochner–Riesz (box multiplier versus disc multiplier, see also Fefferman [12, 13]).

5. Riesz transforms and Poisson integrals. See Stein [22, Chapter III] and [23, Section §4.4].
6. Fourier series and Fourier transform.
  - (a) Basics and summation methods in one and higher dimensions. See Krantz [17, Chapter 1–3], Wolff [29, Chapter 1–5] und Grafakos [14, Chapter 3–4].
  - (b) Mihlin–Hörmander multiplier theorem for Fourier multipliers [24, Chapter VI, Section §4.4]
  - (c) (Optional) Bernstein estimates, and Littlewood–Paley inequalities and their application in nonlinear PDEs.
7. (Optional) Introduction to pseudodifferential operators following Stein [24, Chapter VI] (see also Martinez [18] (Chapter 2, in particular from Section 2.5 on)).
  - (a) Symbolic calculus, composition [24, Chapter VI, Section §3] and [18, Sections 2.6, 2.7].
  - (b)  $L^2$  boundedness, Calderón–Vaillancourt theorem [24, Chapter VI, Section §2] and [18, Section 2.8].
  - (c) Singular integral representation, bounds on integral kernels [24, Chapter VI, Section §4.1-4.3].
  - (d)  $L^2$  boundedness of translation invariant Calderón–Zygmund operators [24, Chapter VI, Section §4.5].
  - (e) Estimates in  $L^p$ , Sobolev, and Lipschitz spaces [24, Chapter VI, Section §5].
8. (Optional) Multiplier theorems for spectral multipliers. Fundamental works by Hebisch [15], Duong–Ouhabaz–Sikora [11], Blunck–Kunstmann [3], and Blunck [2] for Schrödinger and more general self-adjoint operators.
9. (Optional) Almost orthogonality following Stein [24, Chapter VII].
  - (a) Exotic and forbidden symbols, failure of  $L^2$  boundedness for symbols in  $S_{1,1}^0$  [24, Chapter VII, Section §1].
  - (b) Cotlar–Stein lemma [24, Chapter VII, Section §2.1-2.3] and generalization to Schatten classes, see Carbery [7]
  - (c) Consequences of Cotlar–Stein for symbols in  $S_{\rho,\rho}^0$ . (with  $0 \leq \rho < 1$ ) [24, Chapter VII, Section §2.4-2.5].
  - (d)  $L^2$  theory for Calderón–Zygmund operators [24, Chapter VII, Section §3].
  - (e) More on the Cauchy integral [24, Chapter VII, Section §4].
10. (Optional) Oscillatory integrals following Stein [24, Chapter XIII, IX].

- (a) Oscillatory integrals of the first kind, stationary phase [24, Chapter XIII, Section §1-2].
  - (b) Fourier transform of surface measures [24, Chapter XIII, Section §3] and application to the lattice counting problem (improvement of Weyl's law for  $-\Delta$  on  $\mathbb{T}^d$ ) (Sogge [19, pp. 83-85]).
  - (c) Introduction to Fourier restriction [24, Chapter XIII, Section §4].
  - (d) Oscillatory integrals of the second kind, Carleson–Sjölin and Hörmander integral operators [24, Chapter IX, Section §1]. See also Bourgain [4].
  - (e) Relation to Fourier restriction and Bochner–Riesz summability [24, Chapter IX, Section §2], Sogge [20, Sections 2.2-2.3], Tao [26, 27, 28].
11. (Optional) Decoupling inequalities [5, 6, 16].

## References

- [1] S. Blunck and P. C. Kunstmann. Generalized Gaussian estimates and the Legendre transform. *J. Operator Theory*, 53(2):351–365, 2005.
- [2] Sönke Blunck. A Hörmander-type spectral multiplier theorem for operators without heat kernel. *Ann. Sc. Norm. Super. Pisa Cl. Sci. (5)*, 2(3):449–459, 2003.
- [3] Sönke Blunck and Peer Christian Kunstmann. Calderón-Zygmund theory for non-integral operators and the  $H^\infty$  functional calculus. *Rev. Mat. Iberoamericana*, 19(3):919–942, 2003.
- [4] Jean Bourgain. Some new estimates on oscillatory integrals. In *Essays on Fourier Analysis in Honor of Elias M. Stein (Princeton, NJ, 1991)*, volume 42 of *Princeton Math. Ser.*, pages 83–112. Princeton Univ. Press, Princeton, NJ, 1995.
- [5] Jean Bourgain and Ciprian Demeter. The proof of the  $\ell^2$  decoupling conjecture. *Ann. of Math. (2)*, 182(1):351–389, 2015.
- [6] Jean Bourgain and Ciprian Demeter. A study guide for the  $\ell^2$  decoupling theorem. *Chin. Ann. Math. Ser. B*, 38(1):173–200, 2017.
- [7] Anthony Carbery. Almost-orthogonality in the Schatten-von Neumann classes. *J. Operator Theory*, 62(1):151–158, 2009.
- [8] A. Cordoba and R. Fefferman. On the equivalence between the boundedness of certain classes of maximal and multiplier operators in Fourier analysis. *Proc. Nat. Acad. Sci. U.S.A.*, 74(2):423–425, 1977.

- [9] Miguel de Guzmán. *Differentiation of Integrals in  $R^n$* . Lecture Notes in Mathematics, Vol. 481. Springer-Verlag, Berlin-New York, 1975. With appendices by Antonio Córdoba, and Robert Fefferman, and two by Roberto Moriyón.
- [10] Nelson Dunford and Jacob T. Schwartz. *Linear Operators. Part I*. Wiley Classics Library. John Wiley & Sons, Inc., New York, 1988. General theory, With the assistance of William G. Bade and Robert G. Bartle, Reprint of the 1958 original, A Wiley-Interscience Publication.
- [11] Xuan Thinh Duong, El Maati Ouhabaz, and Adam Sikora. Plancherel-type estimates and sharp spectral multipliers. *J. Funct. Anal.*, 196(2):443–485, 2002.
- [12] Charles Fefferman. Inequalities for strongly singular convolution operators. *Acta Math.*, 124:9–36, 1970.
- [13] Charles Fefferman. The multiplier problem for the ball. *Ann. of Math. (2)*, 94:330–336, 1971.
- [14] Loukas Grafakos. *Classical Fourier Analysis*, volume 249 of *Graduate Texts in Mathematics*. Springer, New York, third edition, 2014.
- [15] Waldemar Hebisch. A multiplier theorem for Schrödinger operators. *Colloq. Math.*, 60/61(2):659–664, 1990.
- [16] Jonathan Hickman and Marco Vitturi. Notes on the solution of the  $\ell^2$  decoupling conjecture by Bourgain and Demeter. Available at [https://www.math.sciences.univ-nantes.fr/~vitturi/lecture\\_notes/lecture\\_notes.html](https://www.math.sciences.univ-nantes.fr/~vitturi/lecture_notes/lecture_notes.html), 2015.
- [17] Steven G. Krantz. *A Panorama of Harmonic Analysis*, volume 27 of *Carus Mathematical Monographs*. Mathematical Association of America, Washington, DC, 1999.
- [18] André Martinez. *An Introduction to Semiclassical and Microlocal Analysis*. Universitext. Springer-Verlag, New York, 2002.
- [19] Christopher D. Sogge. *Hangzhou Lectures on Eigenfunctions of the Laplacian*, volume 188 of *Annals of Mathematics Studies*. Princeton University Press, Princeton, NJ, 2014.
- [20] Christopher D. Sogge. *Fourier Integrals in Classical Analysis*, volume 210 of *Cambridge Tracts in Mathematics*. Cambridge University Press, Cambridge, second edition, 2017.
- [21] E. M. Stein. On limits of sequences of operators. *Annals of Mathematics*, 74(1):140–170, 1961.

- [22] Elias M. Stein. *Singular Integrals and Differentiability Properties of Functions*. Princeton Mathematical Series, No. 30. Princeton University Press, Princeton, N.J., 1970.
- [23] Elias M. Stein. *Topics in Harmonic Analysis Related to the Littlewood-Paley Theory*. Annals of Mathematics Studies, No. 63. Princeton University Press, Princeton, N.J.; University of Tokyo Press, Tokyo, 1970.
- [24] Elias M. Stein. *Harmonic Analysis: Real-Variable Methods, Orthogonality, and Oscillatory Integrals*, volume 43 of *Princeton Mathematical Series*. Princeton University Press, Princeton, NJ, 1993. With the assistance of Timothy S. Murphy, Monographs in Harmonic Analysis, III.
- [25] Elias M. Stein and Guido Weiss. *Introduction to Fourier Analysis on Euclidean Spaces*. Princeton University Press, Princeton, New Jersey, 2 edition, 1971.
- [26] Terence Tao. Lecture notes: Restriction theorems and applications. Available at <https://www.math.ucla.edu/~tao/254b.1.99s/>, 1991.
- [27] Terence Tao. From rotating needles to stability of waves: emerging connections between combinatorics, analysis, and PDE. *Notices Amer. Math. Soc.*, 48(3):294–303, 2001.
- [28] Terence Tao. Some recent progress on the restriction conjecture. In *Fourier Analysis and Convexity*, Appl. Numer. Harmon. Anal., pages 217–243. Birkhäuser Boston, Boston, MA, 2004.
- [29] Thomas H. Wolff. *Lectures on Harmonic Analysis*, volume 29 of *University Lecture Series*. American Mathematical Society, Providence, RI, 2003. With a foreword by Charles Fefferman and a preface by Izabella Łaba, Edited by Izabella Łaba and Carol Shubin.